arXiv:math/0201077v5 [math.GN] 15 Feb 2007

A family of pseudo metrics on B^3 and its application

Young Deuk Kim
Department of Mathematics
State University of New York at Stony Brook
Stony Brook, NY 11794-3651, USA
(ydkim@math.sunysb.edu)

February 8, 2008

Abstract

Let B^3 be the closed unit ball in $\mathbf{R}^3$ and S^2 its boundary. We define a family of pseudo metrics on B^3. As an application, we prove that for any countable-to-one function $f : S^2 \to [0, a]$, the set

$$\mathcal{NM}^n_f = \{x \in S^2 \mid \text{there exists } y \in S^2 \text{ such that } f(x) - f(y) > n d_E(x, y)\}$$

is uncountable for all $n \in \mathbf{N}$, where d_E is the Euclidean metric on $\mathbf{R}^3$.

2000 Mathematics Subject Classification ; 57N05, 57M40

1 The family of pseudo metrics

In this section we construct the family of pseudo metrics on the closed unit ball $B^3 \subset \mathbf{R}^3$. As usual, a nonnegative function $d : B^3 \times B^3 \to \mathbf{R}$ is called a *pseudo metric* if

1. $d(x, x) = 0$ for all $x \in B^3$
2. $d(x, y) = d(y, x)$ for all $x, y \in B^3$
3. $d(x, y) + d(y, z) \geq d(x, z)$ for all $x, y, z \in B^3$.

Let $S^2_r \subset \mathbf{R}^3$ be the 2-sphere with center $O = (0, 0, 0)$ and radius $0 < r \leq 1$. We write d_E to denote the Euclidean metric on $\mathbf{R}^3$. A metric d on the set S^2_r is called *locally Euclidean* if for all $P \in S^2_r$, there exists $t > 0$ such that

$$d(Q, R) = d_E(Q, R) \quad \text{for all } Q, R \in B_t(P) = \{S \in S^2_r \mid d(P, S) < t\}.$$

Suppose that $0 < s \leq 1$. Let $-P$ denote the antipodal point of $P \in S^2_r$. Let

$$\alpha = \sin^{-1}\left(\frac{\sqrt{2 - s^2} - s}{2}\right), \quad \text{where } 0 \leq \alpha < \pi/4.$$



Figure 1: S_r^2

Notice that $\sin\alpha$ is a decreasing function of s and hence so is α. For all $P, Q \in S_r^2$, let (see Figure 1)

$$d^s(P,Q) = \begin{cases} d_E(P,Q) & \text{if} \quad \angle POQ \le \pi - 2\alpha \\ 2rs + d_E(-P,Q) & \text{if} \quad \angle POQ > \pi - 2\alpha, \end{cases}$$

where α is defined as above. Notice that if $s = 1$ then $d^1 = d_E$, and $d^s(P,-P) = 2rs$ for all $P \in S_r^2$. In [6] the author proved

Theorem 1.1 *For all* $0 < s \le 1$, d^s *is a locally Euclidean metric on* S_r^2 *which is invariant under any Euclidean isometry.*

Note that (see [6] for a proof)

$$sd_E(P,Q) \le d^s(P,Q) \le d_E(P,Q) \quad \text{for all } P, Q \in S_r^2 \tag{1}$$

and if $d^s(P,Q) \neq d_E(P,Q)$ then

$$d^s(-P,Q) = d_E(-P,Q). \tag{2}$$

We are going to use the metric spaces (S_r^2, d^s) with $0 < r, s \le 1$ to construct a family of pseudo metrics on B^3. Let (see Figure 2)

$$\Delta = \left\{(P,r) \mid P \in int(B^3) \text{ and } 0 < r \le 1 - d_E(O,P)\right\}.$$

For each $\lambda = (P,r) \in \Delta$, choose and fix an isometric embedding ϵ_λ of $\left(S_r^2, d_E\right)$ into (B^3, d_E) such that

$$S_\lambda = \epsilon_\lambda\left(S_r^2\right) = \left\{Q \in B^3 \mid d_E(P,Q) = r\right\}.$$

Suppose that $\Lambda \subset \Delta$ and $P, Q \in B^3$. Let

$$\Pi^n = (B^3)^{n+1} \times \Lambda^n$$
$$[X,\lambda,n] = (X_0, X_1, \cdots, X_n, \lambda_1, \cdots, \lambda_n) \in \Pi^n$$
$$\Gamma_{P,Q} = \{[X,\lambda,n] \in \Pi^n \mid X_0 = P,\ X_n = Q, \text{ and } X_{i-1}, X_i \in S_{\lambda_i} \text{ for all } 1 \le i \le n\}.$$



Figure 2: $\lambda = (P, r) \in \Delta$

Let $\Omega = \{\Lambda \subset \Delta \mid \Gamma_{P,Q} \neq \emptyset$ for all $P, Q \in B^3\}$. Suppose that $\Lambda \in \Omega$ and $s : \Lambda \to (0, 1]$ is a function. Define a function $d^{\Lambda,s} : B^3 \times B^3 \to \mathbf{R}$ by

$$d^{\Lambda,s}(P,Q) = \inf_{[X,\lambda,n]\in\Gamma_{P,Q}} \sum_{i=1}^{n} d^{s(i)}\left({\epsilon_i}^{-1}(X_{i-1}), {\epsilon_i}^{-1}(X_i)\right),$$

where $s(i) = s(\lambda_i)$ and $\epsilon_i = \epsilon_{\lambda_i}$. Notice that $d^{s(i)}\left({\epsilon_i}^{-1}(X_{i-1}), {\epsilon_i}^{-1}(X_i)\right)$ does not depend on ϵ_i because d^s is invariant under Euclidean isometries for all s.

Suppose that $P, Q \in S_\lambda \subset B^3$ for some $\lambda \in \Delta$. Throughout this paper, we write $d^{s(\lambda)}(P,Q)$ to denote $d^{s(\lambda)}\left(\epsilon_\lambda^{-1}(P), \epsilon_\lambda^{-1}(Q)\right)$. Therefore we have

$$d^{\Lambda,s}(P,Q) = \inf_{[X,\lambda,n]\in\Gamma_{P,Q}} \sum_{i=1}^{n} d^{s(i)}(X_{i-1}, X_i).$$

We also write $d_E(P,Q)$ to denote $d_E\left(\epsilon_\lambda^{-1}(P), \epsilon_\lambda^{-1}(Q)\right)$. Notice that there is no problem with this notation because ϵ_λ is an isometric embedding for all $\lambda \in \Delta$.

It is straightforward that $d^{\Lambda,s}$ is a pseudo metric on B^3 for all $\Lambda \in \Omega$ and $s : \Lambda \to (0, 1]$.

2 Elementary properties of the pseudo metric

In this section we study some elementary properties of the pseudo metric. In particular we are interested in the following question.

Question 2.1 *When is $d^{\Lambda,s}$ a metric on B^3? If $d^{\Lambda,s}$ is a metric on B^3, when is $(B^3, d^{\Lambda,s})$ homeomorphic to (B^3, d_E)?*

Suppose that $\Lambda \in \Omega$, $s : \Lambda \to (0, 1]$ and $P, Q \in B^3$. The following subset of $\Gamma_{P,Q}$ will turn out to be useful.

$$\begin{aligned}\Gamma^{\mathcal{A}}_{P,Q} = \big\{&[X, \lambda, n] \in \Gamma_{P,Q} \mid \text{for all } 1 \leq i \leq n, \\ &X_{i-1} \text{ and } X_i \text{ are antipodal on } S_{\lambda_i} \text{ or } \ d^{s(i)}(X_{i-1}, X_i) = d_E(X_{i-1}, X_i)\big\}\end{aligned}$$

Let $[X, \lambda, n] = (X_0, X_1, \cdots, X_n, \lambda_1, \cdots, \lambda_n) \in \Gamma_{P,Q}$ and $\lambda_i = (P_i, r_i)$. Suppose that $d^{s(i)}(X_{i-1}, X_i) \neq d_E(X_{i-1}, X_i)$ for some $1 \leq i \leq n$. Let $-X_{i-1}$ be the antipodal point of X_{i-1} on S_{λ_i}. Notice that

$$\epsilon_i^{-1}(-X_{i-1}) = -\epsilon_i^{-1}(X_{i-1}),$$

where $-\epsilon_i^{-1}(X_{i-1})$ is the antipodal point of $\epsilon_i^{-1}(X_{i-1})$ on $S^2_{r_i}$. From eq. (2) we have $d^{s(i)}(-X_{i-1}, X_i) = d_E(-X_{i-1}, X_i)$. Therefore

$$\begin{aligned}&d^{s(i)}(X_{i-1}, X_i) \\ =\ & 2r_i s(\lambda_i) + d_E(-X_{i-1}, X_i) \\ =\ & d^{s(i)}(X_{i-1}, -X_{i-1}) + d^{s(i)}(-X_{i-1}, X_i).\end{aligned}$$

From this observation we have

$$d^{\Lambda,s}(P, Q) = \inf_{[X,\lambda,n] \in \Gamma^{\mathcal{A}}_{P,Q}} \sum_{i=1}^{n} d^{s(i)}(X_{i-1}, X_i). \tag{3}$$

Suppose that $\Lambda \in \Omega$. Λ is called *piecewise dense* if for all distinct two points $P, Q \in B^3$, there exists $[X, \lambda, n] \in \Gamma_{P,Q}$ (see Figure 3) such that X_i is on the Euclidean segment $\overline{PQ}$ and $d_E(P, X_{i-1}) < d_E(P, X_i)$ for all $1 \leq i \leq n$.



Figure 3: piecewise dense

Note that, for these X_i's, we have $\sum_{i=1}^{n} d_E(X_{i-1}, X_i) = d_E(P, Q)$. Let

$$\Delta_t = \{(P, r) \in \Delta \mid r \leq t\}.$$

Notice that if $\Delta_t \subset \Lambda$ for some $t > 0$ then Λ is piecewise dense.

By the following theorem if Λ is piecewise dense and s is bounded below by some $s_0 > 0$, then $d^{\Lambda,s}$ is a metric on B^3 and the identity map from (B^3, d_E) to $(B^3, d^{\Lambda,s})$ is a homeomorphism.

Theorem 2.1 *Suppose that* $\Lambda \in \Omega$*,* $s : \Lambda \to (0, 1]$ *and* $P, Q \in B^3$*.*

1. *If* Λ *is piecewise dense then* $d^{\Lambda,s}(P, Q) \leq d_E(P, Q)$*.*
2. *If* s *is bounded below by some* $s_0 > 0$ *then* $d^{\Lambda,s}(P, Q) \geq s_0 d_E(P, Q)$*.*

Proof. Suppose that Λ is piecewise dense and $P, Q \in B^3$. We may assume that $P \neq Q$. Choose $[X, \lambda, n] \in \Gamma_{P,Q}$ such that X_i is on the Euclidean segment $\overline{PQ}$ and $d_E(P, X_{i-1}) < d_E(P, X_i)$ for all $1 \leq i \leq n$. From eq. (1) we have

$$d^{\Lambda,s}(P, Q) \leq \sum_{i=1}^{n} d^{s(i)}(X_{i-1}, X_i) \leq \sum_{i=1}^{n} d_E(X_{i-1}, X_i) = d_E(P, Q).$$

Suppose that s is bounded below by some $s_0 > 0$ and $P, Q \in B^3$. From eq. (1) we have

$$\begin{aligned} d^{\Lambda,s}(P, Q) &= \inf_{[X,\lambda,n]\in\Gamma_{P,Q}} \sum_{i=1}^{n} d^{s(i)}(X_{i-1}, X_i) \\ &\geq \inf_{[X,\lambda,n]\in\Gamma_{P,Q}} \sum_{i=1}^{n} s(\lambda_i) d_E(X_{i-1}, X_i) \\ &\geq \inf_{[X,\lambda,n]\in\Gamma_{P,Q}} \sum_{i=1}^{n} s_0 d_E(X_{i-1}, X_i) \\ &= s_0 \inf_{[X,\lambda,n]\in\Gamma_{P,Q}} \sum_{i=1}^{n} d_E(X_{i-1}, X_i) \\ &\geq s_0 d_E(P, Q). \end{aligned}$$

∎

3 The application

The application in this section is motivated by the famous 3-dimensional Poincaré conjecture.

Poincaré Conjecture *If a compact connected 3-manifold is homotopic to* S^3 *then it is homeomorphic to* S^3*.*

See [1, 5, 8, 9, 10, 11] for the recent progress on the Poincaré conjecture by Perelman. The work of Perelman is geometric and analytic. See [2, 3, 4, 7, 12, 13, 14] for the topological approach to the Poincaré conjecture.

Let (M, d) be a compact connected 3-manifold with metric d which is homotopic to S^3. Suppose that $h : (M, d) \to (S^3, d_E)$ be the homotopy equivalence, where d_E is the Euclidean metric on $S^3 \subset \mathbf{R}^4$. Let $g : (S^3, d_E) \to (M, d)$ be any function such that $g \circ h(m) = m$ for all $m \in M$. Note that g is not necessarily continuous.

Question 3.1 *Suppose that* $x, y \in S^3$. *How can we compare* $d(g(x), g(y))$ *with* $d_E(x, y)$*?*

One possible approach to the above question is as follows. For a fixed point $m_0 \in M$, define a function $f : (S^3, d_E) \to \mathbf{R}$ by $f(x) = d(m_0, g(x))$. Notice that $d(g(x), g(y)) \geq f(x) - f(y)$ for all $x, y \in S^3$ and $f(S^3) \subset [0, a]$ for some $a > 0$ because (M, d) is compact. Therefore the following question could be a subquestion of Question 3.1.

Question 3.2 *Suppose that* $f : (S^3, d_E) \to [0, a]$ *is a function and* $x, y \in S^3$. *What can we say about* $f(x) - f(y)$ *and* $d_E(x, y)$*?*

There is no countable-to-one continuous function from (S^3, d_E) to the closed interval $[0, a]$. And any countable-to-one function on S^3 induces countable-to-one functions on subsets of S^3. As an application of the pseudo metric of previous sections, we will prove the following theorem in the next section.

Theorem 3.1 (Main Theorem) *Let* S^2 *be the boundary of* $B^3 \subset \mathbf{R}^3$. *Suppose that* $a > 0$ *and* $f : S^2 \to [0, a]$ *is a countable-to-one function. Then* $\mathcal{NM}_f^n$ *is uncountable for all* $n \in \mathbf{N}$, *where*

$$\mathcal{NM}_f^n = \left\{ x \in S^2 \mid \text{there exists } y \in S^2 \text{ such that } f(x) - f(y) > n d_E(x, y) \right\}.$$

Note that $\mathcal{NM}_f^n$ does not contain the set of discontinuities of f. For example, if C be a countable dense subset of S^2 and

$$f(x) = \begin{cases} 1 & \text{if } x \in C \\ 0 & \text{otherwise} \end{cases}$$

then $\mathcal{NM}_f^n = C$ for all $n \in \mathbf{N}$ but f is discontinuous at every point on S^2.

Notice that Theorem 3.1 is not true for $S^1 = \{(x, y) \mid x^2 + y^2 = 1\} \subset \mathbf{R}^2$. For example, if $g(x) = d_E(x, x_0)$ for some fixed $x_0 \in S^1$ then $g^{-1}(t)$ has at most two elements for all t but $\mathcal{NM}_g^1 = \emptyset$ because

$$g(x) - g(y) = d_E(x, x_0) - d_E(y, x_0) \leq d_E(x, y) \quad \text{for all } x, y \in S^1.$$

Notice also that countability of $f^{-1}(t)$ is crucial in Theorem 3.1. For example, if $h(x) = d_E(x, x_0)$ for some fixed $x_0 \in S^2$ then $h^{-1}(t)$ is uncountable for all $0 < t < 2$ and $\mathcal{NM}_h^1 = \emptyset$.

4 Proof of Theorem 3.1

In this section we prove Theorem 3.1. We write S^2 to denote the boundary of $B^3 \subset \mathbf{R}^3$ throughout this section. First, we will show that the following theorem implies Theorem 3.1.

Theorem 4.1 *Suppose that* $f : S^2 \to [0, 1]$ *is a countable-to-one function. Then* $\mathcal{NM}_f^2$ *is uncountable.*

Theorem 4.1 implies the following theorem.

Theorem 4.2 *Suppose that* $a > 0$ *and* $f : S^2 \to [0, a]$ *is a countable-to-one function. Then* $\mathcal{NM}_f^2$ *is uncountable.*

Proof. Choose $n \in \mathbf{N}$ such that $n > a$ and define a function g by $g(x) = \frac{1}{n} f(x)$. Then $g : S^2 \to \left[0, \frac{a}{n}\right] \subset [0,1]$ and

$$\begin{aligned}
\mathcal{NM}_f^2 &= \{x \in S^2 \mid \exists y \in S^2 \text{ such that } f(x) - f(y) > 2d_E(x,y)\} \\
&= \{x \in S^2 \mid \exists y \in S^2 \text{ such that } ng(x) - ng(y) > 2d_E(x,y)\} \\
&\supset \{x \in S^2 \mid \exists y \in S^2 \text{ such that } g(x) - g(y) > 2d_E(x,y)\} \\
&= \mathcal{NM}_g^2.
\end{aligned}$$

Notice that $g^{-1}(t)$ is countable for all t because $g^{-1}(t) = f^{-1}(nt)$. Therefore by Theorem 4.1, $\mathcal{NM}_g^2$ is uncountable. Hence $\mathcal{NM}_f^2$ is uncountable. ∎

Due to Theorem 4.2, we can prove Theorem 3.1.

Proof of Theorem 3.1. Define a function h by $h(x) = \frac{1}{n} f(x)$. Notice that $\mathcal{NM}_f^n \supset \mathcal{NM}_h^2$. Notice also that $h^{-1}(t)$ is countable for all t because $h^{-1}(t) = f^{-1}(nt)$. Therefore by Theorem 4.2, $\mathcal{NM}_h^2$ is uncountable. Hence $\mathcal{NM}_f^n$ is uncountable. ∎

In the remaining of this paper we will prove Theorem 4.1. Suppose that $f : S^2 \to [0,1]$ is a function. Notice that if $x \in S^2$ then $\frac{x}{2} \in int(B^3)$. From now on, we will use the following fixed $\Lambda \in \Omega$ and $s : \Lambda \to (0,1]$ to prove Theorem 4.1 (see Figure 4).

$$\begin{aligned}
&\Lambda = \Delta_{\frac{1}{2}} = \left\{(P,r) \in \Delta \mid r \leq \frac{1}{2}\right\} \\
&s\left(\frac{x}{2}, \frac{1}{2}\right) = \frac{1 + f(x)}{2} \quad \text{for all } x \in S^2 \quad \text{and} \quad s = 1 \text{ otherwise.}
\end{aligned}$$

Notice that Λ is piecewise dense and the function s is bounded below by $\frac{1}{2}$. By Theorem 2.1, $d^{\Lambda,s}$ is a metric on B^3 and the identity map from (B^3, d_E) to $(B^3, d^{\Lambda,s})$ is homeomorphism.

The following two lemmas will turn out to be useful.

Lemma 4.1 *If* $d^{\Lambda,s}(P,Q) < \frac{1}{2}$ *then* $d^{\Lambda,s}(P,Q) = d_E(P,Q)$ *for all* $P, Q \in B^3$.

Proof. Suppose that $P, Q \in B^3$ and $d^{\Lambda,s}(P,Q) < \frac{1}{2}$. Let

$$\Gamma_{P,Q}^E = \left\{[X, \lambda, n] \in \Gamma_{P,Q}^{\mathcal{A}} \mid \text{for all } 1 \leq i \leq n,\ d^{s(i)}(X_{i-1}, X_i) = d_E(X_{i-1}, X_i)\right\}.$$



Figure 4: $s(P,r)=1$ if $r\neq\frac{1}{2}$ or $S_{(P,r)}$ does not intersect S^2.

Suppose that $[X,\lambda,n]\in\Gamma^{\mathcal{A}}_{P,Q}\setminus\Gamma^{E}_{P,Q}$. There exists j such that X_{j-1} and X_j are antipodal on S_{λ_j} with $\lambda_j=\left(\frac{x}{2},\frac{1}{2}\right)$ for some $x\in S^2$. Therefore

$$\sum_{i=1}^{n} d^{s(i)}(X_{i-1},X_i)\geq d^{s(j)}(X_{j-1},X_j)=\frac{1+f(x)}{2}\geq\frac{1}{2}.$$

Recall that $d^{\Lambda,s}(P,Q)<\frac{1}{2}$. Therefore from eq. (3), we have

$$\begin{aligned}
d^{\Lambda,s}(P,Q) &= \inf_{[X,\lambda,n]\in\Gamma^{\mathcal{A}}_{P,Q}}\sum_{i=1}^{n} d^{s(i)}(X_{i-1},X_i)\\
&= \inf_{[X,\lambda,n]\in\Gamma^{E}_{P,Q}}\sum_{i=1}^{n} d^{s(i)}(X_{i-1},X_i)\\
&= \inf_{[X,\lambda,n]\in\Gamma^{E}_{P,Q}}\sum_{i=1}^{n} d_E(X_{i-1},X_i)\\
&\geq d_E(P,Q).
\end{aligned}$$

Since Λ is piecewise dense, we have $d^{\Lambda,s}(P,Q)\leq d_E(P,Q)$. Therefore

$$d^{\Lambda,s}(P,Q)=d_E(P,Q).$$

■

Lemma 4.2 *If* $d_E(P,Q)<\frac{1}{2}$ *then* $d^{\Lambda,s}(P,Q)=d_E(P,Q)$ *for all* $P,Q\in B^3$.

Proof. Suppose that $d_E(P,Q)<\frac{1}{2}$. Since Λ is piecewise dense, we have $d^{\Lambda,s}(P,Q)\leq d_E(P,Q)<\frac{1}{2}$. Therefore from Lemma 4.1, we have $d^{\Lambda,s}(P,Q)=$

$d_E(P,Q)$. ∎

Suppose that $O=(0,0,0)\in B^3$ and $x\in S^2$. Recall from eq. (3) that

$$d^{\Lambda,s}(O,x)=\inf_{[X,\lambda,n]\in\Gamma^{\mathcal{A}}_{O,x}}\sum_{i=1}^{n} d^{s(i)}(X_{i-1},X_i). \tag{4}$$

The following subsets of $\Gamma^{\mathcal{A}}_{O,x}$ will turn out to be useful.

$$\begin{aligned}\Gamma^{\mathcal{U}}_{O,x} &= \Big\{[X,\lambda,n]\in\Gamma^{\mathcal{A}}_{O,x}\mid \text{there exists unique } i \text{ such that } X_{i-1} \text{ and } X_i \\ &\qquad \text{are antipodal on } S_{\lambda_i} \text{ with } \lambda_i=\left(\frac{y}{2},\frac{1}{2}\right) \text{ for some } y\in S^2\Big\}\\ \Gamma^{\partial}_{O,x} &= \Big\{[X,\lambda,n]\in\Gamma^{\mathcal{U}}_{O,x}\mid \text{there exists } y\in S^2 \text{ such that}\\ &\qquad X_1=y \text{ and } \lambda_1=\left(\frac{y}{2},\frac{1}{2}\right)\Big\}.\end{aligned}$$

Notice that for all $x\in S^2$, we have

$$\left(O,x,\left(\frac{x}{2},\frac{1}{2}\right)\right)\in\Gamma^{\partial}_{O,x}\subset\Gamma^{\mathcal{U}}_{O,x}\subset\Gamma^{\mathcal{A}}_{O,x} \tag{5}$$

$$d^{\Lambda,s}(O,x)\le d^{s\left(\frac{x}{2},\frac{1}{2}\right)}(O,x)=\frac{1+f(x)}{2}\le 1. \tag{6}$$

By the following two lemmas, we can use $\Gamma^{\partial}_{O,x}$ instead of $\Gamma^{\mathcal{A}}_{O,x}$ in eq. (4).

Lemma 4.3 *Suppose that $x\in S^2$. Then*

$$d^{\Lambda,s}(O,x)=\inf_{[X,\lambda,n]\in\Gamma^{\mathcal{U}}_{O,x}}\sum_{i=1}^{n} d^{s(i)}(X_{i-1},X_i).$$

Proof. Suppose that $[X,\lambda,n]\in\Gamma^{\mathcal{A}}_{O,x}\setminus\Gamma^{\mathcal{U}}_{O,x}$. If there is no i such that X_{i-1} and X_i are antipodal on S_{λ_i} with $\lambda_i=\left(\frac{y}{2},\frac{1}{2}\right)$ for some $y\in S^2$, then $d^{s(i)}(X_{i-1},X_i)=d_E(X_{i-1},X_i)$ for all $1\le i\le n$. Therefore

$$\sum_{i=1}^{n} d^{s(i)}(X_{i-1},X_i)=\sum_{i=1}^{n} d_E(X_{i-1},X_i)\ge d_E(O,x)=1.$$

If there exist two distinct integers j,k such that X_{j-1} and X_j are antipodal on S_{λ_j} with $\lambda_j=\left(\frac{y}{2},\frac{1}{2}\right)$, and X_{k-1} and X_k are antipodal on S_{λ_k} with $\lambda_k=\left(\frac{z}{2},\frac{1}{2}\right)$ for some $y,z\in S^2$, then

$$\begin{aligned}\sum_{i=1}^{n} d^{s(i)}(X_{i-1},X_i) &\ge d^{s(j)}(X_{j-1},X_j)+d^{s(k)}(X_{k-1},X_k)\\ &\ge \frac{1+f(y)}{2}+\frac{1+f(z)}{2}\\ &\ge 1.\end{aligned}$$

Therefore by eq. (5) and (6), we have

$$d^{\Lambda,s}(O,x) = \inf_{[X,\lambda,n]\in\Gamma^{\mathcal{U}}_{O,x}} \sum_{i=1}^{n} d^{s(i)}(X_{i-1},X_i).$$

∎

Lemma 4.4 *Suppose that $x \in S^2$. Then*

$$d^{\Lambda,s}(O,x) = \inf_{[X,\lambda,n]\in\Gamma^{\partial}_{O,x}} \sum_{i=1}^{n} d^{s(i)}(X_{i-1},X_i).$$

Proof. Suppose that $[X,\lambda,n] \in \Gamma^{\mathcal{U}}_{O,x} \setminus \Gamma^{\partial}_{O,x}$. There exists unique $j \neq 1$ such that X_{j-1} and X_j are antipodal on S_{λ_j} with $\lambda_j = \left(\frac{y}{2},\frac{1}{2}\right)$ for some $y \in S^2$. Let $X_{j-1} = z$ and $X_j = -z$ (see Figure 5). Note that z and $-z$ are antipodal on S_{λ_j}. Note also that O and y are antipodal on S_{λ_j}, too.

If $d^{\Lambda,s}(O,z) \geq \frac{1}{2}$ then from eq. (5) and (6), we have

$$\begin{aligned} \sum_{i=1}^{n} d^{s(i)}(X_{i-1},X_i) &\geq \sum_{i=1}^{j-1} d^{s(i)}(X_{i-1},X_i) + d^{s\left(\frac{y}{2},\frac{1}{2}\right)}(z,-z) \\ &\geq d^{\Lambda,s}(O,z) + d^{s\left(\frac{y}{2},\frac{1}{2}\right)}(z,-z) \\ &\geq \frac{1}{2} + \frac{1+f(y)}{2} \\ &\geq 1 \\ &\geq \frac{1+f(x)}{2} \\ &\geq \inf_{[X,\lambda,n]\in\Gamma^{\partial}_{O,x}} \sum_{i=1}^{n} d^{s(i)}(X_{i-1},X_i). \qquad (7) \end{aligned}$$

If $d^{\Lambda,s}(O,z) < \frac{1}{2}$ then from Lemma 4.1 and 4.2, we have

$$d^{\Lambda,s}(O,z) = d_E(O,z) = d_E(y,-z) = d^{\Lambda,s}(y,-z).$$

Therefore

$$\begin{aligned} & \sum_{i=1}^{n} d^{s(i)}(X_{i-1},X_i) \\ = \; & \sum_{i=1}^{j-1} d^{s(i)}(X_{i-1},X_i) + d^{s\left(\frac{y}{2},\frac{1}{2}\right)}(z,-z) + \sum_{i=j+1}^{n} d^{s(i)}(X_{i-1},X_i) \\ \geq \; & d^{\Lambda,s}(O,z) + \frac{1+f(y)}{2} + d^{\Lambda,s}(-z,x) \end{aligned}$$



Figure 5: $z \neq O$

$$
\begin{aligned}
&= d^{\Lambda,s}(y,-z) + \frac{1+f(y)}{2} + d^{\Lambda,s}(-z,x) \\
&\geq \frac{1+f(y)}{2} + d^{\Lambda,s}(y,x) \\
&= \frac{1+f(y)}{2} + \inf_{[X,\lambda,n]\in\Gamma_{y,x}} \sum_{i=1}^{n} d^{s(i)}(X_{i-1},X_i). \qquad (8)
\end{aligned}
$$

Suppose that $(Y_0, Y_1, \cdots, Y_m, \lambda'_1, \cdots, \lambda'_m) \in \Gamma_{y,x}$ and let $s'(i) = s(\lambda'_i)$ for all $1 \leq i \leq m$. If there exists j such that Y_{j-1} and Y_j are antipodal on $S_{\lambda'_j}$ with $\lambda'_j = \left(\frac{w}{2}, \frac{1}{2}\right)$ for some $w \in S^2$, then from eq. (5) and (6) we have

$$
\begin{aligned}
\frac{1+f(y)}{2} + \sum_{i=1}^{m} d^{s'(i)}(Y_{i-1},Y_i) \geq \frac{1+f(y)}{2} + d^{s'(j)}(Y_{j-1},Y_j) = \frac{1+f(y)}{2} \\
+ \frac{1+f(w)}{2} \geq 1 \geq \frac{1+f(x)}{2} \geq \inf_{[X,\lambda,n]\in\Gamma^{\partial}_{O,x}} \sum_{i=1}^{n} d^{s(i)}(X_{i-1},X_i).
\end{aligned}
$$

If there is no i such that Y_{i-1} and Y_i are antipodal on $S_{\lambda'_i}$ with $\lambda'_i = \left(\frac{w}{2}, \frac{1}{2}\right)$ for some $w \in S^2$, then from the definition of $\Gamma^{\partial}_{O,x}$ we have

$$
\frac{1+f(y)}{2} + \sum_{i=1}^{m} d^{s'(i)}(Y_{i-1},Y_i) \geq \inf_{[X,\lambda,n]\in\Gamma^{\partial}_{O,x}} \sum_{i=1}^{n} d^{s(i)}(X_{i-1},X_i).
$$

Therefore in eq. (8) we have

$$
\frac{1+f(y)}{2} + \inf_{[X,\lambda,n]\in\Gamma_{y,x}} \sum_{i=1}^{n} d^{s(i)}(X_{i-1},X_i) \geq \inf_{[X,\lambda,n]\in\Gamma^{\partial}_{O,x}} \sum_{i=1}^{n} d^{s(i)}(X_{i-1},X_i).
$$

Therefore from eq. (5), (6), (7) and Lemma 4.3, we have

$$d^{\Lambda,s}(O,x) = \inf_{[X,\lambda,n]\in\Gamma^{\partial}_{O,x}} \sum_{i=1}^{n} d^{s(i)}(X_{i-1},X_i).$$

∎

Due to Lemma 4.4 we can prove the following proposition which will be used in the proof of Theorem 4.1.

Proposition 4.1 *Suppose that* $x \in S^2$*. If* $d^{\Lambda,s}(O,x) \neq \frac{1+f(x)}{2}$ *then* $x \in \mathcal{NM}^2_f$*.*

Proof. Suppose that $x \in S^2$ and $d^{\Lambda,s}(O,x) \neq \frac{1+f(x)}{2}$. From eq. (6) we have

$$d^{\Lambda,s}(O,x) < \frac{1+f(x)}{2}.$$

By Lemma 4.4, there exists $y \in S^2$ such that

$$\left(O, y = X_1, X_2, \cdots, X_{n-1}, x = X_n, \left(\frac{1}{2}y, \frac{1}{2}\right), \lambda_2, \cdots, \lambda_n\right) \in \Gamma^{\partial}_{O,x}$$

$$\frac{1+f(x)}{2} > d^{s\left(\frac{y}{2},\frac{1}{2}\right)}(O,y) + \sum_{i=2}^{n} d^{s(i)}(X_{i-1},X_i).$$

Since

$$\frac{1+f(x)}{2} > d^{s\left(\frac{y}{2},\frac{1}{2}\right)}(O,y) + \sum_{i=2}^{n} d^{s(i)}(X_{i-1},X_i) \geq \frac{1+f(y)}{2} + d^{\Lambda,s}(y,x),$$

we have $1 \geq f(x) - f(y) > 2d^{\Lambda,s}(y,x)$. Therefore from Lemma 4.1. we have $d^{\Lambda,s}(y,x) = d_E(y,x)$. Thus $f(x) - f(y) > 2d_E(x,y)$. Hence $x \in \mathcal{NM}^2_f$. ∎

We need the following fact in the proof of Theorem 4.1.

Fact 4.1 $\mathbf{R}^2 \setminus C$ *is path-connected if* C *is a countable subset of* $\mathbf{R}^2$*. Therefore* $S^2 \setminus C$ *is path-connected if* C *is a countable subset of* S^2*.*

Suppose that $A = \mathbf{R}^2 \setminus C$ and $P, Q \in A$. There are uncountable straight lines in A which contain P and there are uncountable straight lines in A which contain Q. We can choose non-parallel two lines to find a path from P to Q.

Now we can prove Theorem 4.1.

Proof of Theorem 4.1. Suppose that $f : S^2 \to [0,1]$ is a countable-to-one function. To get a contradiction suppose that $\mathcal{NM}^2_f$ is countable. Notice that

$\bigcup_{t\in\mathbf{Q}} f^{-1}(t)$ is countable, where $\mathbf{Q}$ is the set of rational numbers. Therefore the following set A is path-connected by Fact 4.1.

$$A = S^2 \setminus \left(\left(\bigcup_{t\in\mathbf{Q}} f^{-1}(t) \right) \cup \mathcal{NM}_f^2 \right)$$

Define a function $g : (A, d_E) \to [0,1]$ by $g(x) = d^{\Lambda,s}(O,x)$. Since Λ is piecewise dense, we have

$$|g(x) - g(y)| = |d^{\Lambda,s}(O,x) - d^{\Lambda,s}(O,y)| \leq d^{\Lambda,s}(x,y) \leq d_E(x,y).$$

Therefore g is a continuous function and hence $g(A)$ is path-connected.

From Proposition 4.1, we have $g(x) = \frac{1+f(x)}{2}$ for all $x \in A$. Therefore $g(x)$ is an irrational number for all $x \in A$. Notice that $g^{-1}(t)$ is countable for all t and A is a uncountable set. Therefore $g(A)$ is a uncountable subset of irrational numbers. Therefore $g(A)$ is not path-connected. This is a contradiction. Therefore $\mathcal{NM}_f^2$ is uncountable. ∎

References

[1] M. Anderson, *Geometrization of 3-manifolds via the Ricci flow*, Notices Amer. Math. Soc. **51**(2004), no. 2, 184-193.

[2] R. H. Bing, *Mapping a 3-sphere onto a homotopy 3-sphere*, Topology Seminar (Wisconsin, 1965), 89-99, Ann. of Math. Studies, No. 60, Princeton Univ. Press, Princeton, N.J.

[3] D. Gillman and D. Rolfsen, *The Zeeman conjecture for standard spines is equivalent to the Poincaré conjecture*, Topology **22**(1983), no. 3, 315-323.

[4] W. Haken, *On homotopy 3-spheres*, Illinois J. Math. **10**(1966), 159-178.

[5] R. Hamilton, *Three-manifolds with positive Ricci curvature*, J. Differential Geom. **17**(1982), no. 2, 255-306.

[6] Y.D. Kim, *Locally Euclidean metrics on S_r^2 in which some open balls are not connected*, preprint.

[7] J. Milnor, *A unique decomposition theorem for 3-manifolds*, Amer. J. Math. **84**(1962), 1-7.

[8] J. Milnor, *Towards the Poincaré conjecture and the classification of 3-manifolds*, Notices Amer. Math. Soc. **50**(2003), no. 10, 1226-1233.

[9] G. Perelman, *The entropy formula for the Ricci flow and its geometric applications*, preprint, math. DG/0211159.

[10] G. Perelman, *Ricci flow with surgery on three-manifolds*, preprint, math. DG/0303109.

[11] G. Perelman, *Finite extinction time for the solutions to the Ricci flow on certain three-manifolds*, preprint, math. DG/0307245.

[12] V. Poénaru, *A program for the Poincaré conjecture and some of its ramifications*, Topics in low-dimensional topology (University Park, PA, 1996), 65-88, World Sci. Publishing, River Edge, NJ, 1999.

[13] C. Rourke, *Characterisation of the three sphere following Haken*, Turkish J. Math. **18**(1994), no. 1, 60-69.

[14] C. Rourke, *Algorithms to disprove the Poincaré conjecture*, Turkish J. Math. **21**(1997), no. 1, 99-110.